
Bellanita.

A second early, small-sized salad variety with a very high tuberisation and an excellent taste.

- Long tubers with a consistant shape and a nice yellow skin and flesh colour.
- Rather firm cooking with good consumption characteristics.
- Resistant to PCN (RO1&4)* and potato wart disease (1)* and a good resistance to virus and fusarium.



Characteristics

Plant characteristics

Maturity second early Initial development very good Flower colour white

Tuber characteristics

Skin colour yellow Flesh colour yellow Tuber shape long very high **Tuberisation** Tuber size medium Regularity of size regular Superficiality of eyes very shallow Dry matter content 18% Cooking type

Harvest damage slightly sensitive not sensitive Black spot Dormancy short

Resistances

Golden nematode: RO 1 and 4 resistant* Golden nematode: RO 2 and 3 susceptible White nematode: PA 2 susceptible White nematode: PA 3 susceptible Yⁿ-virus immune Common scab quite susceptible Late blight-tuber quite susceptible Late blight-foliage susceptible quite susceptible Spraing Wart disease 1 resistant*

^{*} Results marked with an asterisk (*) are based on our own research, and are not yet officially published.



potato.



Agrico

P.O. Box 70, 8300 AB Emmeloord / The Netherlands Duit 15, 8305 BB Emmeloord / The Netherlands T +31 (0)527 639911 / E info@agrico.nl

I www.agrico.nl



Fertilization¹


Nitrogen: standard advice standard advice Phosphate: Potassium: standard advice

Seed-treatment

Desprouting can be done, but not more than once. Bellanita can be planted directly out of the cold store.

Planting distance

28/35: 22 cm (60,500 plants/ha). 35/50: 25 cm (54,000 plants/ha).





Fungicides

A strict preventative spraying schedule against Late Blight is necessary, as Bellanita is susceptible.



Harvest & Storage



Harvest

Two weeks after accurate haulm killing is usually sufficient, as Bellanita is only a little sensitive to harvest damage and not sensitive to black spot.

Storage

Bellanita has a short dormancy and is only suitable for delivery straight from the field or short term storage.

